

4 février 2008

PAR-LPTHE 01/27
math-ph/0105050**TWO-DIMENSIONAL PARALEL TRANSPORT :
COMBINATORICS AND FUNCTORIALITY**Romain Attal ¹*Laboratoire de Physique Théorique et Hautes Énergies ²
Universités Pierre et Marie Curie (Paris 6) et Denis Diderot (Paris 7)
Unité associée au CNRS UMR 7589*

Abstract : We extend the usual notion of parallel transport along a path to triangulated surfaces. In the one-dimensional case, each path in a base space X is lifted into a fiber bundle with connection and this operation defines a morphism between the fibers above the boundary points of the path. In two dimensions, we define a fibered category over the space $\hat{\Pi}$ of paths in X and we associate to each homotopy of paths a functor between the fibers above the two boundary paths of the swept surface. These "sweeping functors" transport the information from the initial path to the final one. We discuss the abelianisation problem and we conjecture that the smooth limit of the sweeping functors, although not invariant under the action of the group of diffeomorphisms of the swept surface, will provide a space of representation for this group.

PACS : xxxx

¹E-mail : attal@lpthe.jussieu.fr

²LPTHE tour 16 / 1^{er} étage, Université P. et M. Curie, BP 126, 4 place Jussieu, F-75252 Paris Cedex 05 (France)

1 Introduction and physical motivation

In classical and quantum physics, point particles carry information (momentum, spin, color, ...) along their world-line and this evolution is obtained by lifting a path $(\gamma : [0, 1] \rightarrow X)$, described by this particle in the space-time manifold X , into a vector bundle $(E \rightarrow X)$ carrying a representation, R , of a group G . Then a connection on E provides a map $(f_\gamma : F_x \rightarrow F_y)$ between the fibers F_x and F_y above the end points $x = \gamma(0)$ and $y = \gamma(1)$. f_γ is the parallel transport operator associated to all these data. When the path is closed ($x = y$), we obtain an automorphism of the fiber F_x , the trace of which is invariant under the action of G . Since the work of K. Wilson [1], we have a criterion for confinement of quarks in strongly coupled non-Abelian gauge theories : the basic observables are holonomy correlators, $W(\gamma) = \langle \text{tr}(f_\gamma) \cdot \dots \rangle$. The asymptotics $W(\gamma) \sim \exp(-\lambda A_{\min}(\gamma))$, valid in the strongly coupled phase, reflects the collimation of flux lines and the concentration of energy on stringlike excitations sweeping a surface in space-time, with a string tension $\sim \lambda$. Moreover, if we wish to define order/disorder operators in Yang-Mills theory, analogous to those defined in two dimensions, we need a general notion of two-dimensional parallel transport. In order to extend the definition of holonomies to surfaces mapped into a manifold X , we will try to answer the following questions :

- Which fibered spaces are adapted to lift surfaces ?
- How to define a non-Abelian notion of two-dimensional parallel transport ?

It is known that gerbes (a kind of fibered space whose fibers are categories) are the correct setting to answer these questions ([2], [10], [11]), but we do not yet have a full understanding of these spaces when the symmetries form a non-Abelian group, as in gauge theories.

Let G be a locally compact, topological group. This will be our local symmetry group. We will take a polyhedron P as a base space in order to have a locally finite number of degrees of freedom. We have in mind the triangulation of a surface immersed in a manifold. Our building blocks will be the vertices, edges and triangles of this polyhedron. If we want to define a discretized gauge theory from these data, we can associate a copy of G to each vertex and an element of G to each edge, and interpret this as a discretized version of G -bundle with connection. More generally, in order to describe non-trivial bundles, we associate to each vertex a manifold having G as group of automorphisms (the fiber is a G -torsor) and to each edge a morphism between the neighbouring fibers. These data define a functor from the category Π , which has the vertices of P as objects and its edge-paths as arrows, to the category G_1 of right G -torsors. These categories are in fact groupoids since their arrows are isomorphisms. We extend this construction one dimension higher using not only the vertices and edges but also the oriented triangles of P . With these building blocks we define a groupoid $\hat{\Pi}$ whose objects are the edge-paths of P and whose arrows are the simplicial homotopies between these edge-paths. Then we can send $\hat{\Pi}$ in a suitable groupoid $\mathcal{G}$ to represent algebraically our triangulated surfaces. The group G acts on the arrows of $\mathcal{G}$ which are local degrees of freedom associated to the triangles. Then we give a combinatorial definition of the operator which represents our triangulated surface in $\mathcal{G}$. This is not a morphism, as was the case in one dimension, but a functor (between fibered categories) which we call sweeping functor (SF).

In §2, we review some elementary concepts of category theory. We have tried to provide concise definitions and a few examples to be as self-contained as possible. In §3, we introduce the simplicial objects which are the building blocks of the groupoids Π and $\hat{\Pi}$. In §4, we give a definition of simplicial G -bundles. In §5, we extend this definition to fibered categories and we define the sweeping functors as the parallel transport operation in fibered categories. In §6, we illustrate these concepts with a simple example : a 2-sphere triangulated by a tetrahedron and swept in two different ways in order to compare the sweeping functors thus defined. In §7, we discuss the abelianisation problem. A few perspectives are presented in §8.

2 Category vocabulary

2.1 Basic definitions

In this section, we define the concepts of category, groupoid, monoidal category and 2-category [4]. When we compose maps in a category, it is often easier to follow the arrows. Thus we will have $uvw \cdots = \cdots \circ w \circ v \circ u$.

Definitions :

- a) A category $\mathcal{C}$ consists in the following data :
 - a class of objects O ;
 - a class of arrows (or morphisms) A ;
 - two maps $s, t : A \rightarrow O$ (source and target);
 - an associative composition of arrows $((u, v) \mapsto uv)$ defined whenever $t(u) = s(v)$;
 - for each $x \in O$, an identity arrow 1_x which is a left and right neutral element for the composition.
- b) A functor $(f : \mathcal{C} \rightarrow \mathcal{C}')$ is a map which sends each object $x \in O$ to an object $f(x) \in O'$ and each arrow $u \in A$ to an arrow $f(u) \in A'$ such that $f(uv) = f(u)f(v)$ whenever u and v are composable. If f reverses the arrows, i.e. $f(uv) = f(v)f(u)$, it is called a contravariant functor, or simply cofunctor.
- c) A natural transformation $(n : f \rightarrow g)$ between two functors $(f, g : \mathcal{C} \rightarrow \mathcal{C}')$ is a class of arrows $(n_x : f(x) \rightarrow g(x))$, one for each $x \in O$. The naturality of n means that for any arrow $(u : x \rightarrow y)$ of $\mathcal{C}$ we have the commuting square in $\mathcal{C}'$:

$$f(u) n_y = n_x g(u)$$

The composition of arrows in $\mathcal{C}'$ induces a similar composition of natural transformations

$(f \xrightarrow{n} g \xrightarrow{m} h)$ with $n_x m_{f(x)} = (nm)_x \forall x \in O$. The functors $(f : \mathcal{C} \rightarrow \mathcal{C}')$ and their natural transformations form a category noted $\mathcal{F}(\mathcal{C}, \mathcal{C}')$.

d) A functor $f \in \mathcal{F}(\mathcal{C}, \mathcal{C}')$ is an equivalence if there exists another functor $g \in \mathcal{F}(\mathcal{C}', \mathcal{C})$ and two natural transformations $(n : fg \rightarrow \text{Id}_{\mathcal{C}})$ and $(n' : gf \rightarrow \text{Id}_{\mathcal{C}'})$ whose components are isomorphisms.

e) A groupoid is a category where all the arrows are isomorphisms.

f) A category $\mathcal{C}$ is said monoidal if it is equipped with a functor $(\times : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C})$, a unit object I , and natural isomorphisms

$$\begin{aligned} a[A, B, C] &: (AB)C \xrightarrow{\sim} A(BC) \\ \ell_A &: IA \xrightarrow{\sim} A \\ r_A &: AI \xrightarrow{\sim} A \end{aligned}$$

called, respectively, associators, left unit isomorphisms and right unit isomorphisms, and satisfying the pentagon coherence rule

$$\begin{aligned} & \left(((AB)C)D \xrightarrow{a[AB, C, D]} (AB)(CD) \xrightarrow{a[A, B, CD]} A(B(CD)) \right) \\ = & \left(((AB)C)D \xrightarrow{a[A, B, C] \times 1_D} (A(BC))D \xrightarrow{a[A, BC, D]} A((BC)D) \xrightarrow{1_A \times a[B, C, D]} A(B(CD)) \right) \end{aligned}$$

and the triangle coherence rule

$$\left((AI)B \xrightarrow{a[A, I, B]} A(IB) \xrightarrow{1_A \times \ell_B} AB \right) = \left((AI)B \xrightarrow{r_A \times 1_B} AB \right)$$

g) A gr-category is a monoidal groupoid $\mathcal{C}$ where every object A has an inverse A^* : there exists an isomorphism $(\varepsilon_A : A^*A \xrightarrow{\sim} I)$ and this implies the existence of an isomorphism $(\eta_A : AA^* \xrightarrow{\sim} I)$. Roughly speaking, a gr-category is a category endowed with a group-like structure [5][6].

Examples :

i) The finite dimensional vector spaces over a fixed field $\mathbb{F}$ and their $\mathbb{F}$ -linear maps form the category $\text{Vect}_{\mathbb{F}}$.

ii) A topological space T equipped with a continuous, free and transitive left (resp. right) action of a topological group G is called a left (resp. right) G -torsor. A morphism of left (resp. right)

G -torsors is a continuous, G -equivariant map, i.e. $(f : T \rightarrow T')$ satisfies $f(g \cdot t) = g \cdot f(t)$ (resp. $f(t \cdot g) = f(t) \cdot g$) $\forall t \in T$ and $\forall g \in G$. Such maps are in fact homeomorphisms and these data form the category of left (resp. right) G -torsors, denoted G_1^L (resp. G_1^R). We will drop the L or R superscript and consider only right G -torsors.

iii) A mix of the two previous examples is provided by the category of finite dimensional $\mathbb{F}$ -linear representations of G , where the morphisms are the G -equivariant linear maps. This category is denoted $Rep_{\mathbb{F}}(G)$. The tensor product of representation endows $Rep_{\mathbb{F}}(G)$ with a structure of monoidal category.

iv) In § 5, we will use the gr-category G' whose objects are the G -bitorsors. These are the topological spaces on which the group G acts on the left and on the right, freely and transitively. The inverse K^{-1} of a bitorsor K is obtained by exchanging the left and right actions of G . The groupoid G_1 is itself a torsor under the action of G' . Indeed, multiplying on the right any right G -torsor $T \in O(G_1)$ by a bitorsor $K \in O(G')$ defines a right G -torsor TK :

$$\begin{aligned} TK &= \{(t, k) \in T \times K\} / \sim \\ (t, k) &\sim (t \cdot g, g^{-1} \cdot k) \quad \forall g \in G \end{aligned}$$

The inverse operation is the right multiplication by K^{-1} [7].

2.2 2-categories

In a set, the elements are only zero-dimensional objects without internal structure. A category contains also one-dimensional elements (the arrows) which encode the internal structure of its objects. It is natural to extend this notion one step higher to two-dimensions. This is the task of 2-categories.

Definition :

A 2-category $\mathcal{C}$ is defined by the following data :

- a class of objects O ;
- a class of 1-arrows (A_1) and a class of 2-arrows (A_2) ;
- a 1-source map $s_1 : A_2 \rightarrow A_1$ and a 1-target map $t_1 : A_2 \rightarrow A_1$;
- a 0-source map $s_0 : A \rightarrow O$ and a 0-target map $t_0 : A \rightarrow O$ satisfying

$$\begin{aligned} s_0 t_1 &= s_0 s_1 \\ t_0 t_1 &= t_0 s_1 \end{aligned}$$

- a horizontal composition $\times : A_p \times A_p \rightarrow A_p$ defined whenever the 0-source of the first p -arrow is the 0-target of the second one and for $p = 1$ or 2 ;
- a vertical composition $\circ : A_2 \times A_2 \rightarrow A_2$ defined whenever the 1-source of the first 2-arrow is the 1-target of the second one ;
- a left and right identity 1-arrow ($1_x : x \rightarrow x$) for each object x and a left and right identity 2-arrow ($\varepsilon_u : u \rightarrow u$) for each 1-arrow u .

If $x, y \in O$, $A_1(x, y)$ will denote the class of 1-arrows having x as source and y as target, and if $u, v \in A_1(x, y)$, $A_2(u, v)$ will denote the class of 2-arrows having u as 1-source and v as 1-target. To simplify the notations, we will sometimes write $\alpha\beta$ for $\beta \circ \alpha$ and uv for $v \times u$. A 2-arrow $\alpha \in A_2(u, v)$ is invertible when there exists $\beta \in A_2(v, u)$ such that $\alpha\beta = \varepsilon_u$ and $\beta\alpha = \varepsilon_v$. A 1-arrow $u \in A_1(x, y)$ is invertible if there exists $u' \in A_1(y, x)$ and an invertible 2-arrow $\alpha \in A_2(uu', 1_x)$. This notion of invertibility "up to homotopy" is adapted to structures having automorphisms. All these data must satisfy several axioms : associativity (up to homotopy) of the composition of 1-arrows, coherence of these homotopies, ... [4]. When all the 2-arrows of $\mathcal{C}$ are invertible and all its 1-arrows are invertible up to homotopy, we say that $\mathcal{C}$ is a 2-groupoid.

Examples :

- i) The class of all small categories, with the functors as 1-arrows and the natural transformations between these functors as 2-arrows, form the prototype of 2-categories.
- ii) In §6, we will define the 2-groupoid Π_2 whose objects are the vertices of our polyhedron, whose arrows are its edge-paths and whose 2-arrows are equivalence classes of simplicial homotopies sweeping the same oriented 2-cells in (possibly) different orders.

3 Simplicial objects and groupoids

We give here some basic definitions about simplicial objects [8] and we introduce the notion of simplicial groupoid. Our approach to homotopical algebra is very naive but uses groupoids as main tool and functoriality as a guideline.

3.1 Simplicial objects

Simplicial objects are the building blocks of the simplest topological spaces : simplicial complexes. This is the only type of spaces that a computer can handle directly.

Definitions :

- A simplicial complex is a family of sets $K = (K_p)_{p \in \mathbb{N}}$ such that for each $p \geq 1$, K_p is a set of subsets $\sigma \subset K_0$ which contain exactly $(p+1)$ elements and such that any non-empty subset of σ belongs to K_q for some $q \leq p$.
- An element of K_0 is called a vertex and an element of K_p is called a p -simplex. K has pure dimension d if $K_d \neq \emptyset$, $K_p = \emptyset$ for any $p \geq d+1$, and K_0 is the union of the d -simplices of K_d .
- The realization of K , denoted $|K|$, is the set of (barycentric coordinate) functions $(\alpha : K_0 \rightarrow [0, 1])$ whose support, $\text{Supp}(\alpha)$, is a simplex and such that $\sum_{v \in K_0} \alpha(v) = 1$. $|K|$ will be endowed with the weak topology (generated by the inverse images of open sets via the α 's).
- If X is a locally compact topological space, a triangulation of X is a homeomorphism $(f : |K| \rightarrow X)$ and $P = (X, K, f)$ is called a polyhedron.
- The edges of P are the couples (x, y) where $\{x, y\}$ is a 1-simplex of P . The degenerated edges are the couples (x, x) where x is a vertex of P . A finite sequence of edges $((x_i, y_i))_{0 \leq i \leq n}$ is composable if $y_i = x_{i+1}$ for $i = 0, \dots, n-1$.
- For all $q \geq 0$, the q -skeleton of P is $X^q = f(|K_q|)$, where $|K_q| = \{\alpha \in |K|, \text{Supp}(\alpha) \in K_q\}$.

3.2 Simplicial groupoids

We define here the simplicial groupoid of a polyhedron as the category built with the vertices and the edge-paths of this polyhedron.

Definition : Let Π be the groupoid whose objects are the vertices of P and whose arrows are the classes of finite sequences $((x_i, x_{i+1}))_{0 \leq i \leq n-1}$ of composable edges, with respect to the following equivalence relation :

$$(\dots, (x_{i-1}, x), (x, x), (x, x_{i+2}), \dots) \sim (\dots, (x_{i-1}, x), (x, x_{i+2}), \dots)$$


The composition of arrows is the concatenation of sequences. The sequences of the form $((x, x), (x, x), (x, x), \dots)$ are identified to $((x, x))$ which is an identity arrow for this composition. The inverse of $((x_i, x_{i+1}))_{0 \leq i \leq n-1}$ is the reversed path $((x_{n-i}, x_{n-i-1}))_{0 \leq i \leq n-1}$. Π will be called the simplicial groupoid of P .

Definition : Two edge-paths $u = ((x_i, x_{i+1}))_{0 \leq i \leq n-1}$ and $v = ((x'_i, x'_{i+1}))_{0 \leq i \leq n'-1}$ are said X^1 -homotopic if they share the same end points and if there exists a succession of cancellations or insertions of neighbouring opposite edges which transform $((x_i, x_{i+1}))_{0 \leq i \leq n-1}$ into $((x'_i, x'_{i+1}))_{0 \leq i \leq n'-1}$. u and v are said X^2 -homotopic if they share the same end points and if $u^{-1}v$ can be described by a loop which is contractible in X^2 .


Definition : An oriented triangle of P is a q -simplex of P ($q = 0, 1$ or 2) with a couple of marked vertices (x, y) (its 0-source and 0-target) and a couple of marked edge-paths (u, v) (its 1-source and 1-target) joining these vertices, such that :

- (i) If $q = 2$, uv^{-1} forms the boundary of a proper triangle.
- (ii) If $q = 1$ then $u = v = (x, y)$ (flat triangle).
- (iii) If $q = 0$ then $u = v = (x, x) = (y, y)$ (point triangle).


Consequently, the oriented triangles of P are of six kinds, represented by the following pictures :



In these first two cases, the two marked points, a and b , are distinct. The left cell $(\alpha : (ab) \rightarrow (ac, cb))$ is the unique oriented triangle going from the edge $u = (ab)$ to the edge-path $v = (ac, cb)$ and the right cell $(\alpha^* : (ac, cb) \rightarrow (ab))$ is the reversed triangle. We will note (acb) the α -cells and $(acb)^*$ the (reversed) α^* -cells.



In these two other cases, the two marked points are identical ($x = y = c$) and one of the two boundary edge-paths is reduced to the degenerated edge, $u = (cc) = 1_c$, while the other one, $v = (ca, ab, bc)$, describes the whole boundary of the 2-simplex. The cell $(\beta : (cc) \rightarrow (ca, ab, bc))$ is the unique oriented triangle going from the trivial edge (cc) to the edge-path (ca, ab, bc) and $(\beta^* : (ca, ab, bc) \rightarrow (cc))$ is the reversed triangle. We will note (cab) the β -cells and $(cab)^*$ the (reversed) β^* -cells.



The last two cases (degenerated triangles) are the identity arrows and correspond, respectively, to $q = 1$ (left) and $q = 0$ (right).

We now define from Π another groupoid $\widehat{\Pi}$ which is the simplicial groupoid of the space of edge-paths in P . The objects of $\widehat{\Pi}$ are the arrows of Π and the arrows of $\widehat{\Pi}$ are defined by generators and relations.

Definition : The elementary arrows of $\widehat{\Pi}$ are the couples (γ, γ') of edge-paths such that :

- (i) γ and γ' have the same 0-source and 0-target ;
- (ii) there exists an oriented triangle σ such that γ is X^1 -homotopic to $s_1(\sigma)$ and γ' is X^1 -homotopic to $t_1(\sigma)$, both relatively to the subset $\{s_0(\sigma), t_0(\sigma)\}$, i.e. these vertices staying fixed.

In the six cases, the p -source and p -target ($p = 0$ or 1) of an elementary arrow are well defined. Since we are building a groupoid, the elementary arrows must also be invertible. Thus we impose the following relations between our generators :

$$\begin{aligned}
(acb)^* \circ (acb) &= \varepsilon_{ab} \\
(acb) \circ (acb)^* &= \varepsilon_{(ac,cb)} \\
(cabc)^* \circ (cabc) &= \varepsilon_{cc} \\
(cabc) \circ (cabc)^* &= \varepsilon_{(ca,ab,bc)}
\end{aligned}$$

Definition : The arrows of $\widehat{\Pi}$ are the finite sequences $\Gamma = ((\gamma_i, \gamma_{i+1}))_{0 \leq i \leq n-1}$ of elementary arrows sharing the same 0-source and 0-target. The p -source and p -target of Γ are defined by

$$\begin{aligned}
s_0(\Gamma) &= s(\gamma_0) \\
t_0(\Gamma) &= t(\gamma_n) \\
s_1(\Gamma) &= \gamma_0 \\
t_1(\Gamma) &= \gamma_n
\end{aligned}$$

If $\Gamma = ((\gamma_i, \gamma_{i+1}))_{0 \leq i \leq n-1}$ and $\Gamma' = ((\gamma'_i, \gamma'_{i+1}))_{0 \leq i \leq n'-1}$ satisfy $\gamma_n = \gamma'_0$, i.e. $t_1(\Gamma) = s_1(\Gamma')$, their vertical composite is the family $\Gamma\Gamma' = ((\gamma_0, \gamma_1), \dots, (\gamma'_{n'-1}, \gamma'_{n'}))$ obtained by concatenation.

The representations of $\widehat{\Pi}$ will provide us with the right notion of two-dimensional parallel transport. But we first need a simplicial analogue of G -bundles with connection, which we introduce in the next section.

In order to parametrize our edge-paths and simplicial homotopies, we need a simplicial analogue of the interval $[0, 1]$. This rôle is played by the simplicial category Δ whose objects are the finite groupoids Δ_p given by

$$\begin{aligned}
O(\Delta_p) &= \{0, \dots, p-1\} \\
A(i, j) &= \{(i, j)\} \quad \forall \quad i, j \in \{0, \dots, p\}
\end{aligned}$$

The composition of arrows in Δ_p is $(i, j)(j, k) = (i, k)$ and the identity arrows are the (i, i) 's. The arrows of Δ are generated by composition of the shift maps

$$\begin{array}{lll} s_{pk} : \Delta_p & \rightarrow & \Delta_{p+1} \\ i & \mapsto & i \quad (i \leq k) \\ i & \mapsto & i + 1 \quad (i > k) \end{array} \qquad \begin{array}{lll} \tilde{s}_{p\ell} : \Delta_p & \rightarrow & \Delta_{p-1} \\ i & \mapsto & i \quad (i \leq \ell) \\ i & \mapsto & i - 1 \quad (i > \ell) \end{array}$$

A symmetric monoidal structure is defined on Δ by $\Delta_p + \Delta_q = \Delta_{p+q}$ and $I = \Delta_0 = \emptyset$.

$$\begin{aligned} A_\Delta(\Delta_p, \Delta_{p+1}) &= \{s_{pk} / k = 0, \dots, p-1\} \\ A_\Delta(\Delta_p, \Delta_{p-1}) &= \{\tilde{s}_{p\ell} / \ell = 0, \dots, p-1\} \end{aligned}$$

An edge-path in P is now interpreted as a functor $(\gamma : \Delta_p \rightarrow \Pi)$, p being the number of edges of γ or its length. $\widehat{\Pi}$ is therefore the category of functors $(\Delta_p \rightarrow \Pi)$:

$$O(\widehat{\Pi}) = \bigcup_{p \in \mathbb{N}} O(\mathcal{F}(\Delta_p, \Pi))$$

The arrows of $\widehat{\Pi}$ are generated by the natural transformations between these functors (this supposes a common Δ_p as source) and by the homotopies connecting edge-paths of different lengths.

4 Simplicial G -bundles

So far, we have encoded some topological information about the polyhedron P in the simplicial groupoids Π and $\widehat{\Pi}$. In order to compute anything, we need an algebraic representation of these spaces. We define below simplicial fiber bundles with connection over our polyhedron P as representations of Π . We have already defined the groupoid G_1 of (left) G -torsors. The latter are topological manifolds which look like G but without marked point.

Definition : A simplicial G -bundle with connection over the polyhedron P is a functor $(f : \Pi \rightarrow G_1)$ from the simplicial groupoid Π to the groupoid G_1 of G -torsors. The usual definition of a simplicial G -bundle refers to a simplicial object in the category of

G -bundles over some space. These objects come without connection. We have used the same terminology hoping there will be no confusion.

Such a functor $(f : \Pi \rightarrow G_1)$ associates to each vertex $a \in X^0$ a G -torsor f_a and to any edge-path γ a morphism of G -torsors $(f_\gamma : f_a \rightarrow f_b)$. If γ is an edge (ab) , we will write it f_{ab} . When γ is a loop based at a , f_γ is an automorphism of f_a , i.e. an element of G . This is the holonomy of f along γ . From now on, we will drop the epithet "simplicial" which will be understood.

Definition : Let $(f : \Pi \rightarrow G_1)$ be a G -bundle with connection over P and $R = (V, \rho)$ be a F -linear representation of G . The vector bundle with linear connection associated to f and R is the functor f_R defined by

$$\begin{aligned} f_R : \Pi &\rightarrow (G_1 \times \text{Rep}_F(G)) / \sim \\ x &\mapsto f_R(x) = (f_x \times V) / \sim \\ xy &\mapsto f_R(xy) = (f_{xy}, \text{Id}_V) \end{aligned}$$

with the usual equivalence relation :

$$(t \cdot g, v) \sim (t, \rho(g)v)$$

A morphism of G -bundles is a natural transformation $(n : f \rightarrow g)$ between the underlying functors i.e. a morphism of G -torsors $(n_a : f_a \rightarrow g_a)$ for each $a \in X^0$, such that $\forall a, b \in X^0$

$$f_{ab} n_b = n_a g_{ab}$$

Two G -bundles f and g are isomorphic when there exists an invertible natural transformation $(n : f \rightarrow g)$. This implies that f and g have the same holonomies. The automorphisms of a G -bundle f (its invertible natural transformations) form the gauge group

$$\text{Aut}(f) = \mathcal{F}(X^0, G)$$

and the G -bundles over P , together with their natural transformations, form the category $\mathbf{Bun}_G(P) = \mathcal{F}(\Pi, G_1)$.

5 Fibered categories and sweeping functors

We now extend the notion of G -bundle with connection one step higher on the n -category ladder. Our approach can be related to the notion of gerbe as developed in ([9], [10], [11]) and

to that of combinatorial stack [12]. Although less general, the discretized versions are easier to visualize than their sheafified parents.

Definition : A (simplicial) gerbe is a functor $(\varphi : \mathcal{B} \rightarrow \mathcal{G})$ from a base groupoid $\mathcal{B}$ to a groupoid $\mathcal{G}$ whose objects are themselves groupoids. The sections of the stack φ are the functors $(\sigma : \mathcal{B} \rightarrow \varphi(\mathcal{B}))$ such that for all $x, y \in O(\mathcal{B})$, $\sigma(x) \in O(\varphi(x))$ and if $(u : x \rightarrow y)$ is an arrow of $\mathcal{B}$, $(\sigma(u) : \sigma(x) \rightarrow \sigma(y))$ is an arrow of the sub-groupoid $\varphi(\mathcal{B})$.

From now on, we will focus on the following particular case. Let $\mathcal{B} = \Pi$ and let $\mathcal{G} = \{G_1\}$ be the groupoid with G_1 as the only object and its automorphisms as arrows. (These automorphisms are the endofunctors obtained by multiplication on the right by a G -bitorsor $K \in O(G')$). For each edge-path γ , Π_γ denotes the sub-groupoid of Π generated by the vertices and edges of γ , φ_γ is the restriction of φ to Π_γ , and S_γ is the category of sections of φ_γ . We define S as the fibered category over $\widehat{\Pi}$, whose fiber at γ is the category S_γ and whose arrows between the fibers S_γ and $S_{\gamma'}$ above two X^2 -homotopic paths γ and γ' will be our sweeping functors. Since φ_γ and $\varphi_{\gamma'}$ are stacks above γ and γ' , a functor $(\psi : S_\gamma \rightarrow S_{\gamma'})$ must send a section σ of φ_γ to a section $\psi(\sigma)$ of $\varphi_{\gamma'}$ and an arrow $(u : \sigma \rightarrow \tau)$ between sections $\sigma, \tau \in O(S_\gamma)$ to an arrow $(\psi(u) : \psi(\sigma) \rightarrow \psi(\tau))$ between the image sections $\psi(\sigma), \psi(\tau) \in O(\varphi_{\gamma'})$. Since the arrows of $\widehat{\Pi}$ were obtained by generators and relations, it is sufficient to define $\mathcal{F}(S_\gamma, S_{\gamma'})$ for neighbouring paths forming an edge (γ, γ') of $\widehat{\Pi}$. So let $\alpha = (acb)$ be an oriented triangle of P . Let (σ_a, σ_b) be a section of (φ_a, φ_b) , and $(\sigma_{ab} : \varphi_{ab}(\sigma_a) \rightarrow \sigma_b)$ be an arrow between these objects. The image $S_\alpha(\sigma_a, \sigma_b) = (\sigma'_a, \sigma'_c, \sigma'_b)$ is the section of $(\varphi_a, \varphi_b, \varphi_c)$ defined by :

$$\begin{aligned}\sigma'_a &= \sigma_a \\ \sigma'_c &= \varphi_{ac}(\sigma_a) \\ \sigma'_b &= \sigma_b\end{aligned}\tag{1}$$

and these objects are connected by the arrows


$$\sigma'_{ac} = 1_{\sigma_c} : \sigma_c \rightarrow \sigma_c\tag{2}$$

$$\sigma'_{cb} = \sigma_{ab} \circ (\varphi_{acb}(\sigma_a))^{-1} : \varphi_{cb}(\sigma'_c) \rightarrow \sigma'_b\tag{3}$$


The inverse move associates to the arrows (τ_{ac}, τ_{cb}) the arrow $(\tau'_{ab} : \varphi_{ab}(\tau_a) \rightarrow \tau_b)$ defined by

$$\tau'_{ab} = \tau_{cb} \circ \varphi_{cb}(\tau_{ac}) \circ \varphi_{acb}(\tau_a)\tag{4}$$

Thus, to each oriented triangle $\alpha = ((acb) : (ab) \rightarrow (ac, cb))$ corresponds a functor $(S_\alpha : S_{(ab)} \rightarrow S_{(ac, cb)})$, and to its inverse α^* corresponds $S_{\alpha^*} = (S_\alpha)^{-1}$:



Similarly, to each oriented triangle $\beta = ((cab) : (cc) \rightarrow (ca, ab, bc))$ corresponds a functor $(S_\beta : S_{(cc)} \rightarrow S_{(ca, ab, bc)})$, and to its inverse β^* corresponds $S_{\beta^*} = (S_\beta)^{-1}$:




Definition : To an arrow $\Gamma = ((\gamma_i, \gamma_{i+1}))_{0 \leq i \leq n-1}$ of $\widehat{\Pi}$ corresponds the functor $(S_\Gamma : S_{\gamma_0} \rightarrow S_{\gamma_n})$ defined by composition :

$$S_\Gamma = (S_{\gamma_1 \gamma_0^{-1}})(S_{\gamma_2 \gamma_1^{-1}}) \cdots (S_{\gamma_{n-1} \gamma_{n-2}^{-1}})(S_{\gamma_n \gamma_{n-1}^{-1}}) \quad (5)$$

This functor S_Γ will be called the sweeping functor (SF) associated to the homotopy Γ .

Remark : To define a SF we need a succession of edge-paths and this induces an orientation of the underlying 2-simplices. Another sequence of edge-paths sweeping the same surface can define a different orientation of the 2-simplices and thus a different SF, as illustrate the following pictures :




This process measures the curvature of the connection of S around a square of $\widehat{\Pi}$ which projects to a degenerated tetrahedron in P :

$$S_{(ab,bd)} \xrightarrow{S_{acb} \otimes 1} S_{(ac,cb,bd)} \xrightarrow{1 \otimes S_{cbd}^{-1}} S_{(ac,cd)} \xrightarrow{S_{abc} \otimes 1} S_{(ab,bc,cd)} \xrightarrow{1 \otimes S_{bcd}^{-1}} S_{(ab,bd)}$$

6 A basic example

Let's consider as an elementary illustration of the previous definitions the case of a tetrahedron swept in two different ways determined by the following pictures :



We start from the path (ac, cb) with the generic word $(\sigma_{ac}, \sigma_{cb}) = (x, y)$ whose letters form a connection above this initial path, we sweep the surface path after path and we come back to the base path (ac, cb) with a final word to be compared to the initial one. The succession of words in the first case is :

$$\begin{aligned} (ac, cb) &\rightarrow (x, y) \\ (ab) &\rightarrow (xy\varphi_{acb}^{-1}) \\ (ad, db) &\rightarrow (xy\varphi_{acb}^{-1}, \varphi_{adb}) \\ (ad, dc, cb) &\rightarrow (xy\varphi_{acb}^{-1}, \varphi_{adb}, \varphi_{dcb}) \\ (ac, cb) &\rightarrow (xy\varphi_{acb}^{-1}\varphi_{adb}\varphi_{adc}^{-1}, \varphi_{dcb}) \end{aligned}$$

And in the second case, we have :

$$\begin{aligned}
(ac, cb) &\rightarrow (x, y) \\
(ab) &\rightarrow (xy\varphi_{acb}^{-1}) \\
(ad, db) &\rightarrow (xy\varphi_{acb}^{-1}, \varphi_{adb}) \\
(ac, cd, db) &\rightarrow (xy\varphi_{acb}^{-1}, \varphi_{acd}, \varphi_{adb}) \\
(ac, cb) &\rightarrow (xy\varphi_{acb}^{-1}, \varphi_{acd}\varphi_{adb}\varphi_{cdb}^{-1})
\end{aligned}$$

These two sweeping (or pasting) schemes are related by a flat tetrahedron, as described in the previous remark. In order to compare the final words to the initial one, we can make a gauge transformation on the final connection so that we obtain a word of the form (xg_1, yg_2) . If we compare this final word to the initial one, the insertion of the g_i 's defines a notion of 2-holonomy. This operation depends of course on the sweeping scheme. In order to obtain an invariant notion of 2-holonomy, we must lift unparametrized surfaces in a fibered category having "more symmetry".

7 Abelianisation

In the previous sections, we have worked on the space $\widehat{\Pi}$ of paths in P . The sweeping functors defined above depend on the sequence of paths γ_i which sweep the surface in X . How can we get rid of this dependence and obtain invariant sweeping functors which depend only on the swept surface? The pasting theorem [13] states that the composition of a family of composable 2-arrows in any 2-category depends only on the planar oriented graph (or pasting scheme) they define, and not on the order of composition. Each pasting scheme is a discrete flow on the swept surface and the evolution of a path is a succession of local homotopies. Two distant homotopies must be represented algebraically by operations which commute. This is indeed the case when we represent these motions by insertions of letters in a word. This commutativity is a natural consequence of locality, but there is another commutativity which appears if we are not careful with the objects on which the infinitesimal homotopies act. A mistake is to put the same fiber, say G_1 , above each trivial loop and to connect them by the identity functor, believing that such a trivial gerbe on the path space can have interesting holonomies on surfaces since only the 2-cells carry the "fluxes" which are our physical degrees of freedom. But if we suppose that the connective structure is trivial, $\phi_{ab} = \text{Id} : G_1 \rightarrow G_1$ is the trivial G -bitorator for all $a, b \in X^0$. For each oriented triangle (acb) , for all $x, y \in O(\varphi_a)$ and for any arrow $(u : x \rightarrow y)$, the naturality of $(\varphi_{acb} : \varphi_{ab} \rightarrow \varphi_{ac}\varphi_{cb})$ means the following equivariance condition :

$$\varphi_{ab}(u)\varphi_{acb}(y) = \varphi_{acb}(x)\varphi_{ac}\varphi_{cb}(u) \quad (6)$$

The relation (6) implies that for $x = y$, $\varphi_{acb}(x)$ commutes with every $u \in \text{Aut}(x) \simeq G$ thus φ_{acb} lands in $Z(G)$, the center of G . This pitfall is related to the argument used by Eckman and Hilton to prove the commutativity of the second homotopy group of a topological space [14].

In fact, the most general sweeping functors transport the sections of the fibered category φ_u , which lives above the initial path u , to a section of φ_v above the final path v , the edge-paths u and v being X^2 -homotopic. These sections are words of objects. By sweeping each oriented triangle, one inserts a group element in these words or regroups neighbouring letters, as indicated by (1-4). This process depends on the order and orientation of the elementary 2-arrows, as shown in the last remark of the previous section. In a smooth limit, to be precisely defined, we expect the sweeping functors to depend on the parametrisation of the swept surface and to form a representation space for the group of diffeomorphisms of this surface.

8 Conclusion and Perspectives

Let's summarise our work. We have considered a polyhedron P as a base space. We have defined its simplicial groupoid Π as the category which has the vertices as objects and the edge-paths as arrows. Having fixed a locally compact topological group G , we have defined the simplicial G -bundles with connection over P as the functors from Π to the groupoid G_1 of (right) G -torsors. The fibers are associated to the vertices of P and are connected by the morphisms associated to the edges of P . Then, we have extended this definition to two dimensions, with the groupoid $\hat{\Pi}$. The objects of $\hat{\Pi}$ are the edge-paths of P and the arrows of $\hat{\Pi}$ are obtained by sweeping a finite number of oriented triangles in a consistent way from an initial path to a final one (with the same boundary points). A fibered category over $\hat{\Pi}$ is then constructed, which associates to each path the category of sections of a stack in groupoids over P , restricted to this path. The connection of this fibered category of sections associates to each combinatorial homotopy Γ between two edge-paths a functor S_Γ which transports the sections of fibered categories along the surface. On a smooth surface Σ embedded in X , the 2-cells of a pasting diagram define a simplicial approximation of a smooth flow and $\text{Diff}(\Sigma)$ could act on the space of general sweeping functors, thus defining a representation of this group of diffeomorphisms.

Perhaps the main lesson of our approach is that the discretization of geometric, set-theoretic objects leads naturally to a description in terms of categories and higher algebraic structures. This functorial approach seems adapted to gauge theories and string theories. Indeed, when the energy concentrates on thin flux tubes, on loops or on membranes, these new excitations carry information on higher dimensional cells and must be lifted into a suitable fibered n -category. A study of these spaces could shed a new light on the duality relations.

Acknowledgements : I wish to thank Daniel Bennequin for his constant help and encouragement, Lawrence Breen who gave me Ariane's thread of functoriality, and Jean-Luc Brylinski for his critics of my early work. I am also grateful to James Stasheff for a careful reading of the preprint.

Note added : During the publication of the present paper, Lawrence Breen informed me of his joint work with William Messing about the differential geometry of gerbes [16]. They found new identities satisfied by the connection, 2-curvature and 3-curvature, which may have a simple interpretation in our functorial context (work in progress ...).

Références

- [1] K. Wilson, *Confinement of Quarks*, Phys. Rev. D **10** (1974), pp. 2445-2459.
- [2] M. Mackaay and R. Picken, *Holonomy and Parallel Transport for Abelian Gerbes*, math.DG/0007053 v2.
- [3] D. Freed, *Higher Algebraic Structures and Quantization*, Comm. Math. Phys. **159** (1994), pp. 343-398.
- [4] S. Mac Lane, *Categories for the Working Mathematician*, Graduate Texts in Mathematics **5**, 2nd edition, Springer Verlag, 1997.
- [5] A. Joyal and R. Street, *Braided Tensor Categories*, Advances in Math. **102** (1993), pp. 20-78.
- [6] L. Breen, *Théorie de Schreier Supérieure*, Ann. Scient. de l'E.N.S. **25**, pp. 465-514 (1992).
- [7] L. Breen, *Bitorseurs et Cohomologie Non-Abélienne*, in The Grothendieck Festschrift, edited by P. Cartier et al., Progress in Mathematics **86**, pp. 401-476, Birkhäuser (1990).
- [8] J. Peter May, *Simplicial Objects in Algebraic Topology*, University of Chicago Press (1967).
- [9] L. Breen, *On the Classification of 2-Gerbes and 2-Stacks*, Astérisque **225** (1994), Société Mathématique de France.
- [10] J.-L. Brylinski, *Loop Spaces, Characteristic Classes and Geometric Quantization*, Progress in Mathematics **107**, Birkhäuser, 1993.
- [11] J. Giraud, *Cohomologie Non-Abélienne*, Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen **179**, Springer-Verlag, 1971.
- [12] M. M. Kapranov, *Analogies between the Langlands Correspondence and Topological Quantum Field Theory*, in Functional Analysis on the Eve of the 21st Century, edited by S. Gindikin et al., Progress in Mathematics **131**, Birkhäuser, 1995.
- [13] A. J. Power, *A 2-Categorical Pasting Theorem*, J. Algebra **129** (1990), pp. 439-445.

- [14] J. C. Baez and J. Dolan, *Higher Dimensional Algebra and Topological Quantum Field Theory*, J. Math. Phys. **36** (11), Nov. 1995.
- [15] R. Street, *The Algebra of Oriented Simplexes*, J. Pure Appl. Algebra **49** (1987), pp. 283-335.
- [16] L. Breen and W. Messing, *Differential Geometry of Gerbes*, arXiv :math.AG/0106083.